

JOHNSON HOMOMORPHISMS AND ACTIONS OF HIGHER-RANK LATTICES ON RIGHT-ANGLED ARTIN GROUPS

RICHARD D. WADE

ABSTRACT. Let G be a real seimsimple Lie group with no compact factors and finite centre, and let Λ be a lattice in G . Suppose that there exists a homomorphism from Λ to the outer automorphism group of a right-angled Artin group A_Γ with infinite image. We give a strict upper bound to the real rank of G that is determined by the structure of cliques in Γ . An essential tool is the Andreadakis–Johnson filtration of the Torelli subgroup $\mathcal{T}(A_\Gamma)$ of $\text{Aut}(A_\Gamma)$. We answer a question of Day relating to the abelianisation of $\mathcal{T}(A_\Gamma)$, and show that $\mathcal{T}(A_\Gamma)$ and its image in $\text{Out}(A_\Gamma)$ are residually torsion-free nilpotent.

1. INTRODUCTION

Right-angled Artin groups have delivered considerable applications to geometric group theory (two examples are the association with special cube complexes [15] and Bestvina–Brady groups [2]). Another recent area of progress has been the extension of rigidity properties held by irreducible lattices in semisimple Lie groups to involve mapping class groups and automorphism groups of free groups [14, 5, 6]. In this paper we look to what extent this phenomenon extends to the automorphism group of a right-angled Artin group A_Γ , where Γ is the defining graph. If Γ is discrete then A_Γ is the free group F_n , and if Γ is complete then A_Γ is the free abelian group $\mathbb{Z}^n$. One therefore expects traits shared by both $\mathbb{Z}^n$ and F_n to be shared by an arbitrary RAAG. Similarly, one optimistically hopes that properties shared by both $\text{GL}_n(\mathbb{Z})$ and $\text{Out}(F_n)$ will also be shared by $\text{Out}(A_\Gamma)$ for an arbitrary right-angled Artin group. For instance, there is a Nielsen-type generating set of $\text{Out}(A_\Gamma)$ given by the work of Laurence [17] and Servatius [23], and $\text{Out}(A_\Gamma)$ has finite virtual cohomological dimension [8]. $\text{Out}(A_\Gamma)$ is residually finite [9, 21], and for a large class of graphs, $\text{Out}(A_\Gamma)$ satisfies the Tits alternative [9].

Bridson and the author recently showed that any homomorphism from an irreducible lattice Λ in a higher-rank semisimple Lie group to $\text{Out}(F_n)$ has finite image [6]. A direct translation of this result cannot hold for an arbitrary RAAG, as $\mathbb{Z}^n$ is a RAAG and $\text{Out}(\mathbb{Z}^n) = \text{GL}_n(\mathbb{Z})$. However, Margulis’ superrigidity implies that if $\Lambda \rightarrow \text{GL}_n(\mathbb{Z})$ is a map with infinite image, then the real rank of the Lie group containing Λ must be less than or equal to $n - 1$, the real rank of $\text{SL}_n(\mathbb{R})$. The aim of this paper is to show that we may effectively bound the rank of an irreducible lattice acting on A_Γ , and that this bound is determined by obvious copies of $\text{SL}_m(\mathbb{Z})$ in $\text{Out}(A_\Gamma)$. We will describe these copies now. Suppose that Γ' is a subgraph of Γ which is a clique (i.e. any two vertices of Γ' are connected by an edge). A free abelian subgroup of rank m in A_Γ does not imply there exists a copy of $\text{SL}_m(\mathbb{Z})$ in $\text{Out}(A_\Gamma)$, however if the set of vertices adjacent to $v \in \Gamma'$ is independent of v then the natural embedding $\mathbb{Z}^{|\Gamma'|} \rightarrow A_\Gamma$ induces an injection $\text{SL}_{|\Gamma'|}(\mathbb{Z}) \rightarrow \text{Out}(A_\Gamma)$.

Date: 26 July 2011.

2010 Mathematics Subject Classification. 20E36, 20F36 (22E40, 20F14).

Define the SL–dimension of $\text{Out}(A_\Gamma)$, written $d_{SL}(\text{Out}(A_\Gamma))$, to be the number of vertices in the largest such graph Γ' .

Theorem 7.6. *Let G be a real semisimple Lie group with finite centre, no compact factors, and $\text{rank}_{\mathbb{R}} G \geq 2$. Let Λ be an irreducible lattice in G . If $\text{rank}_{\mathbb{R}} G \geq d_{SL}(\text{Out}(A_\Gamma))$, then every homomorphism $f : \Lambda \rightarrow \text{Out}(A_\Gamma)$ has finite image.*

Note that $d_{SL}(\text{GL}_n(\mathbb{Z})) = n$ and $d_{SL}(\text{Out}(F_n)) = 1$. Our previous observation that $\text{Out}(A_\Gamma)$ contains a copy of $SL_m(\mathbb{Z})$ for $m = d_{SL}(\text{Out}(A_\Gamma))$ tells us that the bound on $\text{rank}_{\mathbb{R}} G$ given in Theorem 7.6 is the best that one can provide. The above theorem is deduced from the previously mentioned results for $\text{Out}(F_n)$ and $SL_n(\mathbb{Z})$, combined with the following general algebraic criteria:

Theorem 6.3. *Suppose that $d_{SL}(\text{Out}(A_\Gamma)) = m$, and let $F(\Gamma)$ be the size of a maximal, discrete, full subgraph of A_Γ . Let Λ be a group. Suppose that for each finite index subgroup $\Lambda' \leq \Lambda$, we have:*

- *Every homomorphism $\Lambda' \rightarrow SL_m(\mathbb{Z})$ has finite image,*
- *For all $N \leq F(\Gamma)$, every homomorphism $\Lambda' \rightarrow \text{Out}(F_N)$ has finite image.*

Then every homomorphism $f : \Lambda \rightarrow \text{Out}(A_\Gamma)$ has finite image.

The constant $F(\Gamma)$ can be viewed as the maximal rank of a free subgroup of A_Γ that arises from a subgraph of Γ . In the free group case, the rigidity result for lattices was deduced from a more general result that showed that $\mathbb{Z}$ –averse groups, that is, groups for which no normal subgroup maps onto $\mathbb{Z}$, have no interesting maps to $\text{Out}(F_n)$. To do this, one requires some deep geometric results on the behaviour of *fully irreducible* elements of $\text{Out}(F_n)$ [16, 10, 3], and algebraic results about the structure of the Torelli subgroup of $\text{Out}(F_n)$ (the subgroup consisting of automorphisms that act trivially on $H_1(F_n)$). The results for $\text{Out}(A_\Gamma)$ in [8, 9] come from looking at *projection homomorphisms* which, when Γ is connected, allow us to understand $\text{Out}(A_\Gamma)$ in terms of automorphisms of smaller RAAGs. This allows for inductive arguments. Our approach is to combine the projection homomorphisms alluded to above with algebraic results concerning the structure of the Torelli subgroup of $\text{Out}(A_\Gamma)$. After the background material in the following section, the paper is organised like so: . Section 3 uses Lie methods pioneered by Magnus [20] to study the lower central series of A_Γ ; in particular we study the consecutive quotients $L_c = \gamma_c(A_\Gamma)/\gamma_{c+1}(A_\Gamma)$, and the Lie $\mathbb{Z}$ –algebra L formed by taking the direct sum $L = \bigoplus_{c=1}^{\infty} L_c$, where the Lie bracket is induced by taking commutators in A_Γ . Let $Z(\cdot)$ denote the centre of a group or Lie algebra, $(L)_p$ the tensor product of L with $\mathbb{Z}/p\mathbb{Z}$, and $L^c = L / \bigoplus_{i>c} L_i$. The main technical result concerning L is:

Theorem 3.8. *If $Z(A_\Gamma) = \{1\}$ then $Z(L) = Z((L)_p) = 0$ and $Z(L^c)$ is the image of L_c under the quotient map $L \rightarrow L^c$.*

We obtain this by looking at the enveloping algebra of L , which we call $U(A_\Gamma)$. This has a particularly nice description as a free $\mathbb{Z}$ –module with a basis consisting of positive elements of A_Γ . This allows us to study $Z(L)$ by looking at centralisers of elements in A_Γ , which are well understood.

In Section 4 we describe a central filtration $\mathcal{T}(A_\Gamma) = G_1 \supseteq G_2 \supseteq G_3 \dots$ of $\mathcal{T}(A_\Gamma)$ analogous to the Andreadakis–Johnson filtration of $\mathcal{T}(F_n)$. We have Johnson homomorphisms

$$\tau_c : G_c \rightarrow \text{Hom}(L_1, L_{c+1})$$

for which $\ker \tau_c = G_{c+1}$. Day [11] has shown that $\mathcal{T}(A_\Gamma)$ is finitely generated by a set $\mathcal{M}_\Gamma$, which we describe in Section 2.4. We show that τ_1 maps the elements of $\mathcal{M}_\Gamma$ to linearly independent elements in $\text{Hom}(L_1, L_2)$, therefore $H_1(\mathcal{T}(A_\Gamma))$ is a

free abelian group with a basis given by the image of $\mathcal{M}_\Gamma$. This answers Question 5.4 of [11]. In particular, $\mathcal{M}_\Gamma$ is a minimal generating set of $\mathcal{T}(A_\Gamma)$. The above filtration is separating, so that $\bigcap_{c=1}^\infty G_c = \{1\}$, and the groups L_c are free abelian, so each consecutive quotient G_c/G_{c+1} is free abelian. In the final part of Section 4 we combine Theorem 3.8 with results of Minasyan [21] and Toinet [25] to show that the image $H_1, H_2, H_3, \dots$ of the series $G_1, G_2, G_3, \dots$ in $\text{Out}(A_\Gamma)$ satisfies the same properties. (This roughly follows methods designed by Bass and Lubotsky [1] for studying central series.) In particular, we attain the following result:

Theorem 4.8. *For any graph Γ , the group $\overline{\mathcal{T}}(A_\Gamma)$ is residually torsion-free nilpotent.*

This was also discovered independently by Toinet [25]. This is a key part in the proof of Theorem 6.3; in particular we use it to deal with the inductive step when the underlying graph Γ is disconnected.

A problem arises that projection homomorphisms are not generally surjective; consequently they may raise SL-dimension. This is confronted in Section 5, where we study the image of $\text{Out}(A_\Gamma)$ under a projection map. As such subgroups appear naturally when working with projections, we feel that the methods here may be of independent interest. In particular, we extend the definition of SL-dimension to arbitrary subgroups of $\text{Out}(A_\Gamma)$, and show that if we only study the image of $\text{Out}(A_\Gamma)$ under a projection map, SL-dimension does not increase. This prepares us to complete the proof of Theorem 6.3. We describe applications in Section 7.

The author would like to thank his supervisor Martin Bridson for his encouragement and advice, and Andrew Sale for a series of enthusiastic and helpful conversations.

2. BACKGROUND

Most of this section is standard, however there are some ideas here that are new. In Section 2.1 we extend the usual generating set of $\text{Out}(A_\Gamma)$ to include *extended partial conjugations*, which are products of partial conjugations that conjugate by the same element of A_Γ . We also order the vertices of Γ in a useful way in Sections 2.2 and 2.3, which will be used later in the paper to give a block decomposition of the action of subgroups of $\text{Out}(A_\Gamma)$ on $H_1(A_\Gamma)$.

We should first define A_Γ : Let Γ be a graph with vertex and edge sets $E(\Gamma)$ and $V(\Gamma)$ respectively. If $\iota, \tau : E(\Gamma) \rightarrow V(\Gamma)$ are the maps that take an edge to its initial and terminal vertices, then A_Γ has the presentation:

$$A_\Gamma = \langle v : v \in V(\Gamma) \mid [\iota(e), \tau(e)] : e \in E(\Gamma) \rangle.$$

Throughout we assume that Γ has n vertices labelled $\{v_1, \dots, v_n\}$.

2.1. A generating set of $\text{Aut}(A_\Gamma)$. Laurence [17] proved a conjecture of Servatius [23] that $\text{Aut}(A_\Gamma)$ has a finite generating set consisting of the following automorphisms:

Graph symmetries: If a permutation of the vertices comes from a self-isomorphism of the graph, then this permutation induces an automorphism of A_Γ . These automorphisms form a finite subgroup of $\text{Aut}(A_\Gamma)$ called $\text{Sym}(A_\Gamma)$.

Inversions: These are automorphisms that come from inverting one of the generators of A_Γ , so that:

$$s_i(v_k) = \begin{cases} v_i^{-1} & i = k \\ v_k & i \neq k. \end{cases}$$

Partial conjugations: Suppose $[v_i, v_j] \neq 0$. Let Γ_{ij} be the connected component of $\Gamma - st(v_j)$ containing v_i . Then the partial conjugation K_{ij} conjugates every vertex of Γ_{ij} by v_j , and fixes the remaining vertices, so that:

$$K_{ij}(v_k) = \begin{cases} v_j v_k v_j^{-1} & v_k \in \Gamma_{ij} \\ v_k & v_k \notin \Gamma_{ij}. \end{cases}$$

Note that if $lk(v_i) \subset st(v_j)$ then $\Gamma_{ij} = \{v_i\}$, so in this case K_{ij} fixes every basis element except v_i .

Transvections: If $lk(v_i) \subset st(v_j)$, then there is an automorphism ρ_{ij} which acts on the generators of A_Γ as follows:

$$\rho_{ij}(v_k) = \begin{cases} v_i v_j & i = k \\ v_k & i \neq k. \end{cases}$$

There are two important finite index normal subgroups of $\text{Aut}(A_\Gamma)$ that we obtain from this classification. The first is the subgroup generated by inversions, partial conjugations, and transvections and is denoted $\text{Aut}^0(A_\Gamma)$. The second is the smaller subgroup generated by only partial conjugations and transvections. Denote this group $\text{SAut}^0(A_\Gamma)$. In some cases we will need to look at groups generated by (outer) automorphisms that conjugate more than one component of $\Gamma - st(v_j)$ by v_j . Let T be a subset of Γ_{ij} such that no two vertices of T lie in the same connected component of Γ_{ij} . We define an *extended partial conjugation* to be an automorphism of the form $\prod_{t \in T} K_{tj}$. We will abuse notation by describing the images of the above elements in $\text{Out}(A_\Gamma)$ by the same names, so that the groups $\text{Out}^0(A_\Gamma)$ and $\text{SOut}^0(A_\Gamma)$ are defined in the same manner. If $\phi \in \text{Aut}(A_\Gamma)$, we use $[\phi]$ to denote the equivalence class of ϕ in $\text{Out}(A_\Gamma)$. Let $\mathcal{A}_\Gamma$ be the enlarged generating set of $\text{Out}(A_\Gamma)$ given by graph symmetries, inversions, extended partial conjugations, and transvections. We shall be studying subgroups of $\text{Out}(A_\Gamma)$ generated by subsets of $\mathcal{A}_\Gamma$, however some of these groups are not generated by subsets of the standard generating set. This is because under the restriction, exclusion and projection maps defined in Section 2.6, partial conjugations are not always mapped to partial conjugations, but are always mapped to extended partial conjugations. Throughout this paper, we will assume $\text{Aut}(A_\Gamma)$ and $\text{Out}(A_\Gamma)$ act on A_Γ on the left.

2.2. Ordering the vertices of Γ . Given a vertex v , the *link of v* is the set of vertices of Γ adjacent to v . The *star of v* is the union of v and the link of v . We write $lk(v)$ for the link of v , and $st(v)$ for the star of v . Extending the definition of the link and star of a vertex, given any full subgraph Γ' of Γ , the subgraph $lk(\Gamma')$ is defined to be the intersection of the links of the vertices of Γ' , and we define $st(\Gamma') = \Gamma' \cup lk(\Gamma')$. Given any full subgraph Γ' , the right-angled Artin group $A_{\Gamma'}$ injects into A_Γ , so can be viewed as a subgroup.

We may define a binary relation on the vertices by letting $u \leq v$ if $lk(u) \subset st(v)$. This was introduced by Charney and Vogtmann in [8], who showed that the relation is transitive as well as symmetric, so defines a preorder on the vertices. This induces an equivalence relation by letting $u \sim v$ if $u \leq v$ and $v \leq u$. Let $[v]$ denote the equivalence class of the vertex v . We will abuse notation by also using $[v]$ to denote the full subgraph of Γ spanned by the vertices in this equivalence class. The preorder descends to a partial order of the equivalence classes. We say that $[v]$ is *maximal* if it is maximal with respect to this ordering. The vertices of $[v]$ are all either at distance one from each other in Γ , or generate a free subgroup of Γ , therefore $A_{[v]}$ is either a free abelian, or a free non-abelian subgroup of A_Γ . We say that the equivalence class $[v]$ is *abelian* or *non-abelian* respectively. Suppose that there are r equivalence classes of vertices in Γ . We may choose an enumeration of the vertices

so that there exists $1 = m_1 < m_2 < \dots < m_r < n$ such that the equivalence classes are the sets $\{v_1 = v_{m_1}, \dots, v_{m_2-1}\}, \dots, \{v_{m_r}, \dots, v_n\}$. With further rearrangement we may assume that $v_{m_i} \leq v_{m_j}$ only if $i \leq j$. We formally define $m_{r+1} = n + 1$ so that for all i , the equivalence class of $[v_{m_i}]$ contains $m_{i+1} - m_i$ vertices.

2.3. G-ordering vertices. Given a subgroup $G \leq \text{Out}(A_\Gamma)$, we may define the relation $\leq_G$ on the vertices of Γ , by letting $v_i \leq_G v_j$ if either $i = j$ or the element $[\rho_{ij}]$ lies in G . Note that this is a subset of the previous relation $\leq$ defined on the vertices. Also, $\leq_G$ is reflexive by definition and transitive as $\rho_{il} = \rho_{jl}^{-1} \rho_{ij}^{-1} \rho_{jl} \rho_{ij}$, so $\leq_G$ is a preorder, induces an equivalence relation $\sim_G$ on the vertices, and induces a partial ordering of the equivalence classes of $\sim_G$. Let $[v_i]_G$ be the equivalence class of the vertex v_i . Each equivalence class $[v_i]_G$ is a subset of the equivalence class $[v_i]$. In particular the subgroup $A_{[v_i]_G}$ is either free abelian or free and non-abelian, so $[v_i]_G$ may also be described as *abelian* or *non-abelian*. Suppose that there are $r' \geq r$ equivalence classes of vertices in $\sim_G$. We may further refine the enumeration of the vertices given previously so that there exists $1 = l_1 < l_2 < \dots < l_{r'} < n$ such that the equivalence classes of $\sim_G$ are the sets $\{v_1 = v_{l_1}, \dots, v_{l_2-1}\}, \dots, \{v_{l_{r'}}, \dots, v_n\}$, and $v_{l_i} \leq_G v_{l_j}$ only if $i \leq j$. Define $l_{r'+1} = n + 1$ so that for all i , the equivalence class of $[v_{l_i}]_G$ contains $l_{i+1} - l_i$ vertices.

2.4. The Torelli subgroups of $\text{Aut}(A_\Gamma)$ and $\text{Out}(A_\Gamma)$. The abelianisation of A_Γ , denoted $A_\Gamma^{ab} = A_\Gamma/[A_\Gamma, A_\Gamma]$ or $H_1(A_\Gamma)$, is a free abelian group generated by the images of the vertices under the abelianisation map $A_\Gamma \rightarrow A_\Gamma/[A_\Gamma, A_\Gamma]$. This induces a natural map

$$\Phi : \text{Aut}(A_\Gamma) \rightarrow \text{Aut}(A_\Gamma^{ab}).$$

Once and for all we fix the basis of A_Γ^{ab} to be the image of $V(\Gamma)$ under the abelianisation map. This allows us to identify $\text{Aut}(A_\Gamma^{ab})$ with $\text{GL}_n(\mathbb{Z})$. This is the viewpoint which we will take for the rest of the paper. We say that $\ker \Phi = \mathcal{T}(A_\Gamma)$ is the *Torelli subgroup* of $\text{Aut}(A_\Gamma)$. Every partial conjugation lies in $\mathcal{T}(A_\Gamma)$, and if i, j, k are distinct, $v_i \leq v_j, v_k$, and $[v_j, v_k] \neq 0$, then the automorphism

$$K_{ijk}(v_l) = \begin{cases} v_i[v_j, v_k] & l = i \\ v_l & l \neq i, \end{cases}$$

lies in $\mathcal{T}(A_\Gamma)$. The conditions on i, j, k ensure that K_{ijk} is well-defined and non-trivial. In [11], Day proves the following theorem:

Theorem 2.1. *The set of partial conjugations and automorphisms of the form K_{ijk} (where i, j, k are distinct, $v_i \leq v_j, v_k$, and $[v_j, v_k] \neq 0$) generates $\mathcal{T}(A_\Gamma)$.*

As $K_{ijk} = K_{ikj}^{-1}$, we may restrict the generating set to contain K_{ijk} only when $j < k$. We call this generating set $\mathcal{M}_\Gamma$.

As Φ sends $\text{Inn}(A_\Gamma)$ to the identity, we can factor out $\text{Inn}(A_\Gamma)$ to obtain a map

$$\overline{\Phi} : \text{Out}(A_\Gamma) \rightarrow \text{GL}_n(\mathbb{Z}).$$

We define $\overline{\mathcal{T}}(A_\Gamma) = \ker \overline{\Phi}$, and call this group the *Torelli subgroup* of $\text{Out}(A_\Gamma)$.

2.5. Words in A_Γ . At times we will need to discuss words and word length in A_Γ . The support, $\text{supp}(w)$, of a word w on $V(\Gamma) \cup V(\Gamma)^{-1}$ is the full subgraph of Γ spanned by the generators (or their inverses) occurring in w . We say that a word w representing $g \in A_\Gamma$ is *reduced* if there is no subword of the form $v^{\pm 1} w' v^{\mp 1}$ with $\text{supp}(w') \subset \text{st}(v)$. We may pass between two reduced words by repeatedly switching consecutive letters that commute (see [7]) — it follows that we may define $\text{supp}(g)$ to be the support of any reduced word representing g . Similarly the *length* of g is the length of any reduced word representing g . We say that g is *positive* if it has a representative that is a positive word in $V(\Gamma)$.

2.6. Restriction, exclusion, and projection homomorphisms. Suppose that Γ' is a full subgraph of Γ , so that $A_{\Gamma'}$ can be viewed as a subgroup of A_{Γ} in the natural way. Suppose that the conjugacy class of $A_{\Gamma'}$ is preserved by a subgroup $G < \text{Out}(A_{\Gamma})$. Then there is a natural *restriction map* $R_{\Gamma'} : G \rightarrow \text{Out}(A_{\Gamma'})$ obtained by taking a representative of an element $[\phi] \in G$ that preserves $A_{\Gamma'}$. This is well defined as the normaliser of $A_{\Gamma'}$ is of the form $C_{A_{\Gamma}}(A_{\Gamma'}) \cdot A_{\Gamma'}$. Hence if $gA_{\Gamma'}g^{-1} = A_{\Gamma'}$, the action of g on $A_{\Gamma'}$ can be realised by an inner automorphism of $A_{\Gamma'}$. Similarly, if the normal subgroup of A_{Γ} generated by $A_{\Gamma'}$ is preserved by some $G < \text{Out}(A_{\Gamma})$, then there is a natural *exclusion map* $E_{\Gamma'} : G \rightarrow \text{Out}(A_{\Gamma}/\langle\langle A_{\Gamma'} \rangle\rangle) \cong \text{Out}(A_{\Gamma-\Gamma'})$. There are two key examples:

Example 2.2. If Γ is connected and v is a maximal vertex then the conjugacy classes of $A_{[v]}$ and $A_{st[v]}$ are preserved by $\text{Out}^0(A_{\Gamma})$ ([8], Proposition 3.2). Therefore there is a restriction map

$$R_v : \text{Out}^0(A_{\Gamma}) \rightarrow \text{Out}^0(A_{st[v]}),$$

an exclusion map

$$E_v : \text{Out}^0(A_{\Gamma}) \rightarrow \text{Out}^0(A_{\Gamma-[v]}),$$

and a projection map

$$P_v : \text{Out}^0(A_{\Gamma}) \rightarrow \text{Out}^0(A_{lk[v]})$$

obtained by combining the restriction and exclusion maps. We can take the direct sum of these projection maps over all maximal equivalence classes $[v]$ to obtain the *amalgamated projection homomorphism*:

$$P : \text{Out}^0(A_{\Gamma}) \rightarrow \bigoplus_{[v] \text{ maximal}} \text{Out}^0(A_{lk[v]})$$

Example 2.3. If Γ is not connected, then there exists a finite set $\{\Gamma_i\}_{i=1}^k$ of connected graphs containing at least two vertices and an integer N such that $A_{\Gamma} \cong F_N *_{i=1}^k A_{\Gamma_i}$. By looking at the action of the generating set of $\text{Out}^0(A_{\Gamma})$, we find that the conjugacy class of A_{Γ_i} is fixed by $\text{Out}^0(A_{\Gamma})$, therefore for each i we obtain a restriction map

$$R_i : \text{Out}^0(A_{\Gamma}) \rightarrow \text{Out}^0(A_{\Gamma_i}),$$

and as the normal subgroup generated by $*_{i \in I} A_{\Gamma_i}$ is preserved by $\text{Out}(A_{\Gamma})$ there is an exclusion map

$$E : \text{Out}(A_{\Gamma}) \rightarrow \text{Out}(F_N).$$

Charney and Vogtmann have shown that when Γ is connected, the maps in Example 2.2 describe $\text{Out}(A_{\Gamma})$ almost completely. There are two cases: when the centre of A_{Γ} , which we write as $Z(A_{\Gamma})$, is trivial, and when $Z(A_{\Gamma})$ is nontrivial. In the first case, they show the following:

Theorem 2.4 ([8], Theorem 4.2). *If Γ is connected and $Z(A_{\Gamma})$ is trivial, then $\ker P$ is a finitely generated free abelian group.*

In [9], Theorem 2.4 is extended by giving an explicit generating set of $\ker P$, however we will not need this description in the work that follows. When $Z(A_{\Gamma})$ is nontrivial there is a unique maximal abelian equivalence class $[v]$ consisting of the vertices in $Z(A_{\Gamma})$, and we are in the following situation:

Proposition 2.5 ([8], Proposition 4.4). *If $Z(A_{\Gamma}) = A_{[v]}$ is nontrivial, then*

$$\text{Out}(A_{\Gamma}) \cong \text{Tr} \rtimes (GL(A_{[v]}) \times \text{Out}(A_{lk[v]})),$$

where Tr is the free abelian group generated by the transvections $[\rho_{ij}]$ such that $v_i \in lk[v]$ and $v_j \in [v]$. The map to $GL(A_{[v]})$ is given by the restriction map R_v , and the map to $\text{Out}(A_{lk[v]})$ is given by the projection map P_v . The subgroup Tr is the kernel of the product map $R_v \times P_v$.

In the above proposition we do not need to restrict R_v and P_v to $\text{Out}^0(A_\Gamma)$, as every automorphism of A_Γ preserves $Z(A_\Gamma) = A_{[v]}$. When Γ is disconnected, the restriction and exclusion maps of Example 2.3 give us less information. As above, we may amalgamate the restriction maps R_i and the exclusion map E , however in this situation the kernel of the amalgamated map is not abelian — it is the semidirect product of a subgroup of $\overline{\mathcal{T}}(A_\Gamma)$ and the abelian group generated by transvections $[\rho_{ij}]$, where v_i is an isolated vertex of Γ , and v_j is contained in a nontrivial connected component of Γ .

3. THE LOWER CENTRAL SERIES OF A_Γ

In this section we shall gather some results on the lower central series of A_Γ and its associated Lie algebra that we require in the rest of the paper. In Proposition 3.6 we give a basis for the free abelian group $\gamma_2(A_\Gamma)/\gamma_3(A_\Gamma)$ and in Theorem 3.8 we give information about the structure of the Lie algebra $L = \oplus_{i=1}^{\infty} \gamma_i(A_\Gamma)/\gamma_{i+1}(A_\Gamma)$. From Magnus' work, most of the results are well known in the free group case (see, for example, Chapter 5 of [20] or Chapter 2 of [4]), however a little care is required generalising the results to all right-angled Artin groups.

3.1. The Lie algebra L and its enveloping algebra $U(A_\Gamma)$. Many facts about the lower central series of the free group were generalised to right-angled Artin groups by Duchamp and Krob [12, 13]; below we summarise some key points from their work. In the above papers right-angled Artin groups are described as *free partially commutative groups*. Note that the algebra $U(A_\Gamma)$ defined below coincides with the *free partially commutative $\mathbb{Z}$ -algebra* used in Duchamp and Krob's work, as we can get between two positive words in $V(\Gamma)$ representing the same element of A_Γ with a sequence of positive words where consecutive elements differ only by permuting a pair of commuting letters (this says that the *free partially commutative monoid* embeds in A_Γ in a natural way.)

Let $\gamma_c(A_\Gamma)$ be the c th term in the lower central series of A_Γ , so that $\gamma_1(A_\Gamma) = A_\Gamma$ and $\gamma_{c+1}(A_\Gamma) = [\gamma_c(A_\Gamma), A_\Gamma]$. Let $L_c = \gamma_c(A_\Gamma)/\gamma_{c+1}(A_\Gamma)$. For all Γ and c , the group L_c is free abelian. Let $L = \oplus_{i=1}^{\infty} L_i$. As $[\gamma_c(A_\Gamma), \gamma_d(A_\Gamma)] \subset \gamma_{c+d}(A_\Gamma)$, the $\mathbb{Z}$ -module L inherits a graded $\mathbb{Z}$ -Lie algebra structure by taking commutators in A_Γ , and is generated by the images of $v_1, \dots, v_n$ in L_1 . Furthermore L is a free $\mathbb{Z}$ -module, and admits a basis consisting of elements of the form

$$(1) \quad \beta(v_{i_1}, v_{i_2}, \dots, v_{i_k}, v_{i_{k+1}}),$$

where β is a bracket of degree k . (For example, if $k = 2$ then we have a basis consisting of elements either of the form $[x, [y, z]]$ or of the form $[[x, y], z]$.)

Definition 3.1. Let $U(A_\Gamma)$ be the free $\mathbb{Z}$ -module with a basis given by positive elements of A_Γ . Let $U_i(A_\Gamma)$ be the submodule of $U(A_\Gamma)$ spanned by positive elements of A_Γ of length i . Then $U(A_\Gamma) = \oplus_{i=0}^{\infty} U_i(A_\Gamma)$, and multiplication in A_Γ gives $U(A_\Gamma)$ the structure of a graded $\mathbb{Z}$ -algebra.

We will distinguish elements of $U(A_\Gamma)$ from A_Γ by writing positive words in $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ rather than $\{v_1, \dots, v_n\}$. Let $\mathcal{L}(U(A_\Gamma))$ be $U(A_\Gamma)$ endowed with the Lie bracket defined by $[u_1, u_2] = u_1 \cdot u_2 - u_2 \cdot u_1$. Then $U(A_\Gamma)$ and L are related by the following theorem:

Theorem 3.2 ([12] Corollary I.2. and [13] Theorem 2.1). *The map $L_1 \rightarrow U_1(A_\Gamma)$ given by $v_i \gamma_2(A_\Gamma) \mapsto \mathbf{v}_i$ induces an injective Lie algebra homomorphism*

$$\epsilon : L \rightarrow \mathcal{L}(U(A_\Gamma)).$$

Furthermore, this map is graded, so that $L_i \rightarrow U_i(A_\Gamma)$, and has the universal property exhibiting the fact that $U(A_\Gamma)$ is the enveloping algebra of L .

Let $U^\infty(A_\Gamma)$ be the algebra extending $U(A_\Gamma)$ by allowing infinitely many coefficients of a sequence of positive words to be non-zero. Any element of $U^\infty(A_\Gamma)$ can be written uniquely as a power series $a = \sum_{i=0}^\infty a_i$, where a_i is an element of $U_i(A_\Gamma)$. We say that a_i is the *homogeneous part* of a of degree i . Each a_i is a linear sum of positive elements of length i , so is of the form $a_i = \sum_{k=1}^m \lambda_k g_k$, where g_k is a positive element in A_Γ of length i and $\lambda_k \in \mathbb{Z} \setminus \{0\}$. Define $\text{supp}(a_i) = \cup_{k=1}^m \text{supp}(g_k)$. Let $U^*(A_\Gamma)$ be the group of units in $U^\infty(A_\Gamma)$. If a is of the form $a = 1 + \sum_{i=1}^\infty a_i$, then $a \in U^*(A_\Gamma)$ and

$$a^{-1} = 1 - (a_1 + a_2 + \cdots) + (a_1 + a_2 + \cdots)^2 - \cdots = 1 + \sum_{i=1}^\infty c_i.$$

Here $c_1 = -a_1$ and $c_j = -\sum_{i=0}^{j-1} c_i a_{j-i}$ recursively. We have abused notation slightly in the above by writing 1 as the leading coefficient rather than 1.1_{A_Γ} . We may use $U^\infty(A_\Gamma)$ to study A_Γ via the following proposition:

Proposition 3.3. *The mapping $v_i \mapsto 1 + \mathbf{v}_i$ induces a homomorphism $\mu : A_\Gamma \rightarrow U^*(A_\Gamma)$.*

Proof. The mapping $v_i \mapsto 1 + \mathbf{v}_i$ induces a homomorphism $\bar{\mu} : F(V) \rightarrow U^*(A_\Gamma)$. The relations in the standard presentation of A_Γ are sent to the identity in $U^*(A_\Gamma)$, so this descends to a homomorphism $\mu : A_\Gamma \rightarrow U^*(A_\Gamma)$. $\square$

μ is sometimes called the *Magnus map* or *Magnus morphism*. It is important because of the following result of Droms, Duchamp and Krob [13, Theorem 1.2]. Its proof follows the method used by Magnus in the case of the free group, which is covered on page 310 of [20].

Proposition 3.4. *The homomorphism $\mu : A_\Gamma \rightarrow U^*(A_\Gamma)$ is injective.*

There is a central series $M_1 \geq M_2 \geq M_3 \dots$ of $U^*(A_\Gamma)$ defined by letting $a \in M_c$ if and only if $a_i = 0$ when $0 < i < c$. It is related to the lower central series of A_Γ by the following theorem:

Theorem 3.5 ([13], Theorem 2.2). *For all c we have $\mu^{-1}(M_c) = \gamma_c(A_\Gamma)$.*

As $\cap_{c=1}^\infty M_c = \{1\}$, it follows that $\cap_{c=1}^\infty \gamma_c(A_\Gamma) = \{1\}$. By induction on c one can show that for $a, b \in U^*(A_\Gamma)$, the element ab^{-1} lies in M_c if and only if $a_i = b_i$ for $i < c$. This gives one an effective way of studying the lower central series of A_Γ .

3.2. More information on the structure of L . We use μ to find free generating sets for the groups L_c :

Proposition 3.6. *The free abelian group $L_2 = \gamma_2(A_\Gamma)/\gamma_3(A_\Gamma)$ has a basis given by the set $S = \{[v_i, v_j]\gamma_3(A_\Gamma) : i < j, [v_i, v_j] \neq 0\}$.*

Proof. As $\gamma_2(A_\Gamma) = [A_\Gamma, A_\Gamma]$ is the normal closure of S , any element of $g \in \gamma_2(A_\Gamma)$ is of the form $g = g_1 s_1^{e_1} g_1^{-1} \cdots g_k s_k^{e_k} g_k^{-1}$, where $s_i \in S$ and $g_i \in A_\Gamma$. However, $[g_i, s_i] \in \gamma_3(A_\Gamma)$ for all i , therefore the image of g in $\gamma_2(A_\Gamma)/\gamma_3(A_\Gamma)$ is equal to $s_1^{e_1} \cdots s_k^{e_k}$, so L_2 is generated by S . As μ is injective and $\mu^{-1}(M_c) = \gamma_c(A_\Gamma)$ for all c , the map μ induces an injection $\bar{\mu} : \gamma_2(A_\Gamma)/\gamma_3(A_\Gamma) \rightarrow M_2/M_3$, and this can be composed further with the homomorphism $f : M_2/M_3 \rightarrow U_2(A_\Gamma)$ given by $a + M_3 \mapsto a_2$. The free abelian group (or free $\mathbb{Z}$ -module) $U_2(A_\Gamma)$ has a basis consisting of elements of length 2 in A_Γ , the set:

$$\{\mathbf{v}_i^2 : v_i \in V(\Gamma)\} \cup \{\mathbf{v}_i \mathbf{v}_j : v_i, v_j \in V(\Gamma) \text{ and } i < j \text{ or } [v_i, v_j] \neq 0\}.$$

Then $f\bar{\mu}([v_i, v_j]\gamma_3(A_\Gamma)) = \mathbf{v}_i \mathbf{v}_j - \mathbf{v}_j \mathbf{v}_i$, so the images of the elements of S are linearly independent in $U_2(A_\Gamma)$, and form a basis of L_2 . $\square$

We shall use Proposition 3.6 in Section 4 to describe the abelianisation of $\mathcal{T}(A_\Gamma)$. The last thing we need is to use this associative algebra to give us information about the structure of L .

Proposition 3.7. *Let $v \in V(\Gamma)$. Let $a = \sum_{i=1}^{\infty} a_i$ be an element of $U(A_\Gamma)$, and let $b = [\mathbf{v}, a] = \sum_{i=1}^{\infty} b_i$. Then $b_{i+1} = 0$ only if $\text{supp}(a_i) \subset \text{st}(v)$.*

Proof. Suppose that $b_{i+1} = 0$. Let $a_i = \sum_{k=1}^m \lambda_k g_k$ be a decomposition of a_i into a sum of distinct positive elements g_k , where each $\lambda_k \in \mathbb{Z} \setminus \{0\}$. As $b_{i+1} = \mathbf{v}a_i - a_i\mathbf{v} = 0$, we have $\mathbf{v}a_i = a_i\mathbf{v}$. We have two injective maps $g_k \mapsto \mathbf{v}g_k$ and $g_k \mapsto g_k\mathbf{v}$ from the set of positive elements of length i to the set of positive elements of length $i+1$, therefore as $\mathbf{v}a_i = a_i\mathbf{v}$ we have $\{g_k\mathbf{v}\}_{k=1}^m = \{\mathbf{v}g_k\}_{k=1}^m$. Suppose that each g_k has a reduced representative where the first j letters commute with v . For each g_k there exists l such that $g_k\mathbf{v} = \mathbf{v}g_l$, so that $g_k = \mathbf{v}g_l\mathbf{v}^{-1}$. There is a reduced representative w_l of g_l where the first j letters commute with v and we can find a reduced representative of g_k by a sequence of swaps of consecutive commuting letters and one cancellation in vw_lv^{-1} . If v^{-1} is cancelled with a letter v from the last $i-j$ letters of w_l then certainly the first $j+1$ letters in this representative of g_k commute with v . If v^{-1} is cancelled by the initial letter v or one of the first j letters of w_l , then v^{-1} must commute with the last $i-j$ letters of w_l , so every letter in this reduced representative of g_k commutes with v . In either case there is a representative of g_k where the first $j+1$ letters commute with v . By induction this must hold for all $j \leq i$, hence $\text{supp}(g_k) \subset \text{st}(v)$ for all k and $\text{supp}(a_i) \subset \text{st}(v)$ $\square$

The above lets us control the centre of L and other associated structures. Let p be a positive integer and let $(L)_p$ be the Lie algebra obtained by taking the tensor product of $\mathbb{Z}/p\mathbb{Z}$ with L . Let L^c be the quotient algebra $L/\oplus_{i>c} L_i$. We use $Z(-)$ to denote the centre of a Lie algebra. (cf. [4], Chapter 2, Exercise 3.3)

Theorem 3.8. *If $Z(A_\Gamma) = \{1\}$ then $Z(L) = Z((L)_p) = 0$ and $Z(L^c)$ is the image of L_c under the quotient map $L \rightarrow L^c$.*

Proof. Elements of L of the form in Equation (1) map to homogeneous polynomials in $U(A_\Gamma)$ under the map $\epsilon: L \rightarrow \mathcal{L}(U(A_\Gamma))$, where the coefficient of each element of A_Γ is ± 1 . As such elements form a basis of L , it follows that tensoring L with $\mathbb{Z}/p\mathbb{Z}$ corresponds to taking the image of L under the map ϵ and tensoring $U(A_\Gamma)$ with $\mathbb{Z}/p\mathbb{Z}$. Also, as ϵ maps $L_i \rightarrow U_i(A_\Gamma)$, the algebra L^c is isomorphic to the image of L in $U^c = U(A_\Gamma)/\oplus_{i>c} U_i(A_\Gamma)$. If we take a nonzero element $a \in U(A_\Gamma)$ then as $Z(A_\Gamma) = \{1\}$, for each a_i there exists $v \in V(\Gamma)$ such that $\text{supp}(a_i) \not\subset \text{st}(v)$. Then $[\mathbf{v}, a] \neq 0$ by Proposition 3.7, and if a is nonzero in $\mathbb{Z}/p\mathbb{Z} \otimes U(A_\Gamma)$, then so is $[\mathbf{v}, a]$ by looking at coefficients. Similarly if a is nonzero in U^c , there exists $i \leq c$ with $a_i \neq 0$. By picking v such that $\text{supp}(a_i) \not\subset \text{st}(v)$, the $(i+1)$ st homogenous part of $[\mathbf{v}, a]$ is nonzero by Proposition 3.7. The proof then follows by noting that if $a \in \text{im } \epsilon$ then as $\mathbf{v} \in \text{im } \epsilon$, so is $[\mathbf{v}, a]$ for all $v \in V(\Gamma)$. $\square$

4. THE ANDREADAKIS-JOHNSON FILTRATION OF $\mathcal{T}(A_\Gamma)$

In this section we follow the methods of Bass and Lubotsky [1] to extend the notion of *higher Johnson homomorphisms* from the free group setting to general right-angled Artin groups. Coupled with the work in the previous sections of the paper, these allow us to describe the abelianisation of $\mathcal{T}(A_\Gamma)$, and show that $\mathcal{T}(A_\Gamma)$ has a separating central series $G_1, G_2, G_3, \dots$ where each quotient G_i/G_{i+1} is a finitely generated free abelian group. This was first studied in the case of free groups by Andreadakis. We show that the image of this series in $\text{Out}(A_\Gamma)$ satisfies the same results.

4.1. Definition and application to $H_1(\mathcal{T}(A_\Gamma))$. As $\gamma_c(A_\Gamma)$ is characteristic, there is a natural map $\text{Aut}(A_\Gamma) \rightarrow \text{Aut}(A_\Gamma/\gamma_c(A_\Gamma))$. Let G_{c-1} be the kernel of this map. Then $G_0 = \text{Aut}(A_\Gamma)$ and $G_1 = \mathcal{T}(A_\Gamma)$. The following proposition is proved in the same way as Proposition 2.2 of [6].

Proposition 4.1. *Let $\phi \in G_c$, where $c \geq 1$. The mapping $\bar{g} \mapsto \phi(g)g^{-1} \cdot \gamma_{c+2}(A_\Gamma)$ induces a homomorphism*

$$\tau_c: G_c \rightarrow \text{Hom}(L_1, L_{c+1})$$

such that $\ker(\tau_c) = G_{c+1}$.

As $\cap_{c=1}^\infty \gamma_c(A_\Gamma) = \{1\}$, it follows that $\cap_{c=1}^\infty G_c = \{1\}$. As L_1 and L_{c+1} are free abelian, $\text{Hom}(L_1, L_{c+1})$ is free abelian, and therefore G_c/G_{c+1} is free abelian. One can check that $[G_c, G_d] \subset G_{c+d}$, so that $G_1, G_2, G_3, \dots$ is a central series of $\mathcal{T}(A_\Gamma)$. The rank of each L_c has been calculated in [13], although more work is needed to calculate the ranks of the quotients G_c/G_{c+1} . In the free group case G_1/G_2 , G_2/G_3 and G_3/G_4 are known [22, 24] but as yet there is no general formula. In this paper we restrict ourselves to studying the abelianisation of $\mathcal{T}(A_\Gamma)$, using the generating set $\mathcal{M}_\Gamma$ defined in Section 2.4.

Theorem 4.2. *The first Johnson homomorphism τ_1 maps $\mathcal{M}_\Gamma$ to a free generating set of a subgroup of $\text{Hom}(L_1, L_2)$. The abelianisation of $\mathcal{T}(A_\Gamma)$ is isomorphic to the free abelian group on the set $\mathcal{M}_\Gamma$, and G_2 is the commutator subgroup of $\mathcal{T}(A_\Gamma)$.*

Proof. By Proposition 3.6, the free abelian group L_2 has a basis $S = \{[v_i, v_j]\gamma_3(A_\Gamma) : i < j, [v_i, v_j] \neq 0\}$. This allows us to obtain an explicit description of the images of elements of $\mathcal{M}_\Gamma$:

$$\begin{aligned} \tau_1(K_{ij})(v_l) &= \begin{cases} 1\gamma_3(A_\Gamma) & \text{if } v_l \notin \Gamma_{ij} \\ [v_j, v_l]\gamma_3(A_\Gamma) & \text{if } v_l \in \Gamma_{ij} \end{cases} \\ \tau_1(K_{ijk})(v_l) &= \begin{cases} 1\gamma_3(A_\Gamma) & \text{if } l \neq i \\ [v_j, v_k]\gamma_3(A_\Gamma) & \text{if } l = i \end{cases} \end{aligned}$$

These elements are linearly independent in $\text{Hom}(L_1, L_2)$. The second statement follows immediately, and the third follows as $G_2 = \ker(\tau_1)$. $\square$

Corollary 4.3. *$\mathcal{M}_\Gamma$ is a minimal generating set of $\mathcal{T}(A_\Gamma)$.*

4.2. Example: the pentagon. Suppose that Γ is the pentagon shown in Figure 1. In this case, if $v_i \leq v_j$ then $v_i = v_j$, therefore no elements of the form K_{ijk} exist in




FIGURE 1.

$\mathcal{T}(A_\Gamma)$. Removing $st(v_i)$ from Γ leaves exactly one connected component consisting of the two vertices opposite v_i , therefore

$$\mathcal{M}_\Gamma = \{K_{13}, K_{24}, K_{35}, K_{41}, K_{52}\}.$$

Hence $H_1(\mathcal{T}(A_\Gamma)) = G_1/G_2 \cong \mathbb{Z}^5$. Also, $\{[v_1, v_3], [v_1, v_4], [v_2, v_4], [v_2, v_5], [v_3, v_5]\}$ is a set of coset representatives of $\gamma_2(A_\Gamma)$ in $\gamma_3(A_\Gamma)$, therefore $\text{Hom}(L_1, L_2) \cong \mathbb{Z}^{25}$. In particular τ_1 is not surjective (in contrast to the free group situation — see [22]).

4.3. Extension to $\text{Out}(A_\Gamma)$. We'd now like to move our attention to the images of $G_1, G_2, \dots$ in $\text{Out}(A_\Gamma)$, which we will label $H_1, H_2, \dots$ respectively. Let π be the natural projection $\text{Aut}(A_\Gamma) \rightarrow \text{Out}(A_\Gamma)$. The action of an element of A_Γ on itself by conjugation induces a homomorphism $ad: A_\Gamma \rightarrow \text{Aut}(A_\Gamma)$. For each $g \in A_\Gamma \setminus \{1\}$ there exists a unique integer d such that $g \in \gamma_d(A_\Gamma)$ and $g \notin \gamma_{d+1}(A_\Gamma)$. In the language of Section 3, we identify g with the element $g\gamma_{d+1}(A_\Gamma)$ in the submodule L_d of the Lie algebra L . We use this to study the map ad in the following lemma:

Lemma 4.4. *If $Z(A_\Gamma) = \{1\}$, then $g \in \gamma_c(A_\Gamma)$ if and only if $ad(g) \in G_c$.*

Proof. If $g \in \gamma_c(A_\Gamma)$ then $ghg^{-1} = h \pmod{\gamma_{c+1}(A_\Gamma)}$, for all $h \in A_\Gamma$. Hence $ad(g) \in G_c$. Conversely, suppose that $g \notin \gamma_c(A_\Gamma)$. Then $g \in L_d$ for some $d < c$, and by Theorem 3.8, the image of g under the quotient map $L \rightarrow L^c$ is not central. As L is generated by $v_1, \dots, v_n$, there exists v_i such that the image of $[g, v_i]$ is nonzero in L^c . Hence $[g, v_i] \neq 1 \pmod{\gamma_{c+1}(A_\Gamma)}$, so $gv_i g^{-1} \neq v_i \pmod{\gamma_{c+1}(A_\Gamma)}$, and $ad(g) \notin G_c$. $\square$

By the above, $ad(\gamma_c(A_\Gamma)) \subset G_c$, and the induced map $\gamma_c(A_\Gamma)/\gamma_{c+1}(A_\Gamma) \rightarrow G_c/G_{c+1}$ is injective. Furthermore, it is clear that $\pi(ad(A_\Gamma)) = \{1\}$, and if $\phi \in G_c$ but $[\phi] \in H_{c+1}$, then there exists $g \in A_\Gamma$ such that $\phi G_{c+1} = ad(g)G_{c+1}$. As $\phi \in G_c$, by Lemma 4.4 we have $g \in \gamma_c(A_\Gamma)$, hence ϕG_{c+1} is in the image of $\gamma_c(A_\Gamma)/\gamma_{c+1}(A_\Gamma)$. Therefore, when $Z(A_\Gamma) = \{1\}$ we have an exact sequence of abelian groups:

$$0 \rightarrow \gamma_c(A_\Gamma)/\gamma_{c+1}(A_\Gamma) \xrightarrow{\alpha} G_c/G_{c+1} \xrightarrow{\beta} H_c/H_{c+1} \rightarrow 0,$$

where α and β are induced by ad and π respectively.

Theorem 4.5. *If $Z(A_\Gamma) = \{1\}$ and $c \geq 1$ then the group H_c/H_{c+1} is free abelian.*

Proof. By the exact sequence above,

$$H_c/H_{c+1} \cong (G_c/G_{c+1})/(ad(\gamma_c(A_\Gamma))G_{c+1}/G_{c+1}).$$

As G_c/G_{c+1} is free abelian, it is sufficient to show that if there exists $\phi \in G_c$, $g \in A_\Gamma$, and a prime p such that $ad(g)G_{c+1} = \phi^p G_{c+1}$, then there exists $h \in A_\Gamma$ such that $ad(h)G_{c+1} = \phi G_{c+1}$.

Suppose that ϕ , g , and p exist as above. As $\phi \in G_c$, for every $x \in A_\Gamma$, there exists $w_x \in \gamma_{c+1}(A_\Gamma)$ such that $\phi(x) = xw_x$. One can check using commutator calculus that

$$\phi(x)^p = xw_x^p \pmod{\gamma_{c+2}(A_\Gamma)},$$

therefore

$$gxg^{-1} = xw_x^p \pmod{\gamma_{c+2}(A_\Gamma)}.$$

It follows that the image of $[g, x]$ is zero in $(L)_p$, for all $x \in A_\Gamma$, hence the image of g lies in $Z((L)_p)$. By Theorem 3.8, g must lie in the kernel of the map $L \rightarrow (L)_p$, so there exists $h \in \gamma_c(A_\Gamma)$ such that $g = h^p \pmod{\gamma_{c+1}(A_\Gamma)}$. Then $[g, x] = [h^p, x] \pmod{\gamma_{c+2}(A_\Gamma)}$ for all $x \in A_\Gamma$, so $ad(h)^p G_{c+1} = ad(g)G_{c+1} = \phi^p G_{c+1}$. As G_c/G_{c+1} is torsion-free it has unique roots. Hence $ad(h)G_{c+1} = \phi G_{c+1}$. $\square$

We now adapt a well-known fact about $\text{Out}(F_n)$ to $\text{Out}(A_\Gamma)$.

Proposition 4.6. *The intersection $\cap_{c=1}^\infty H_c$ is trivial.*

Proof. Let $\phi \in \text{Aut}(A_\Gamma)$, and suppose that $[\phi] \in H_c$ for all c . Then for every element $g \in A_\Gamma$, we know that $\phi(g)$ is conjugate to g in $A_\Gamma/\gamma_c(A_\Gamma)$ for all c . Toinet has shown that RAAGs are conjugacy separable in finite p -group, (and therefore nilpotent) quotients [25]; this tells us that $\phi(g)$ is conjugate to g . Furthermore, Minasyan ([21], Proposition 6.9) has shown that if ϕ takes every element of A_Γ to a conjugate, then ϕ itself is an inner automorphism. Hence $\cap_{c=1}^\infty H_c = \{1\}$. $\square$

Therefore if $Z(A_\Gamma)$ is trivial then $H_1, H_2, H_3 \dots$ is central series of $\overline{\mathcal{T}}(A_\Gamma)$, with trivial intersection, such that the consecutive quotients H_c/H_{c+1} are free abelian.

Corollary 4.7. *If $Z(A_\Gamma)$ is trivial then $\overline{\mathcal{T}}(A_\Gamma)$ is residually torsion-free nilpotent.*

4.4. The situation when $Z(A_\Gamma)$ is nontrivial. Suppose that $Z(A_\Gamma)$ is nontrivial. Let $[v]$ be the unique maximal equivalence class of vertices in Γ . By Proposition 2.5, there is a restriction map R_v and a projection map P_v like so:

$$\begin{aligned} R_v: \text{Out}(A_\Gamma) &\rightarrow \text{GL}(Z(A_\Gamma)) \cong \text{GL}(A_{[v]}) \\ P_v: \text{Out}(A_\Gamma) &\rightarrow \text{Out}(A_\Gamma/Z(A_\Gamma)) \cong \text{Out}(A_{lk[v]}). \end{aligned}$$

The kernel of the map $R_v \times P_v$ is the free abelian subgroup Tr generated by the transvections $[\rho_{ij}]$ such that $v_i \in lk[v]$ and $v_j \in [v]$. Elements of $\text{Out}(A_\Gamma)$ that lie in Tr act nontrivially on A_Γ^{ab} , as do elements that are nontrivial under R_v . It follows that $\overline{\mathcal{T}}(A_\Gamma)$ is mapped isomorphically under P_v onto $\overline{\mathcal{T}}(A_{lk[v]})$. As the centre of $lk[v]$ is trivial, this lets us promote the above work to any right-angled Artin group:

Theorem 4.8. *For any graph Γ , the group $\overline{\mathcal{T}}(A_\Gamma)$ is residually torsion-free nilpotent.*

5. SL-DIMENSION AND PROJECTION HOMOMORPHISMS FOR SUBGROUPS OF $\text{Out}(A_\Gamma)$.

In Section 6 we will restrict the actions of certain groups on RAAGs of bounded SL-dimension by using an induction argument based on the number of vertices in Γ , combined with projection homomorphisms. Unfortunately our definition of SL-dimension described in the introduction can behave badly under projection, restriction, and inclusion homomorphisms. In particular, if v is a maximal vertex in a connected graph Γ , then it is not always true that $d_{SL}(\text{Out}(A_{lk[v]})) \leq d_{SL}(\text{Out}(A_\Gamma))$. To get round this problem, we will extend the definition of SL-dimension to arbitrary subgroups of $\text{Out}(A_\Gamma)$, and show that if instead we look at the image of $\text{Out}(A_\Gamma)$ under such homomorphisms, then the SL-dimension will not increase (for instance, it will always be the case that $d_{SL}(P_v(\text{Out}(A_\Gamma))) \leq d_{SL}(\text{Out}(A_\Gamma))$). For the remainder of this section we fix a subgroup G of $\text{Out}(A_\Gamma)$ generated by a subset $T \subset A_\Gamma$. We shall assume that T is maximal, so that $T = A_\Gamma \cap G$. We are going to study the action of G on $H_1(A_\Gamma)$, so to simplify matters we will often want to ignore graph symmetries and inversions.

Proposition 5.1. *Let G^0 be the subgroup of G generated by the extended partial conjugations, inversions, and transvections in T and let SG^0 be the subgroup of G generated solely by the extended partial conjugations and transvections in T . Then G^0 and SG^0 are finite index normal subgroups of G .*

Proof. Suppose α is a graph symmetry that moves the vertices according to the permutation σ . We find that $\alpha K_{ij} \alpha^{-1} = K_{\sigma(i)\sigma(j)}$, $\alpha \rho_{ij} \alpha^{-1} = \rho_{\sigma(i)\sigma(j)}$ and $\alpha s_i \alpha^{-1} = s_{\sigma(i)}$. As we assumed that T is maximal, if $[\phi]$ and $[\alpha]$ belong to T , then so does $[\alpha\phi\alpha^{-1}]$. Therefore if W is a word in $T \cup T^{-1}$ one may shuffle graph symmetries along so that they all occur at the beginning of W . As the group of graph symmetries is finite, this shows that G^0 is finite index in G , and the above

computations verify that G^0 is normal in G . Similarly, with inversions one verifies that:

$$\rho_{kl}s_i = \begin{cases} s_i\rho_{kl} & i \neq k, l \\ s_i\rho_{ki}^{-1} & i = l \\ s_iK_{il}^{-1}\rho_{il}^{-1} & i = k, [v_i, v_l] \neq 0 \\ s_i\rho_{il}^{-1} & i = k, [v_i, v_l] = 0 \end{cases} \text{ and } K_{kl}s_i = \begin{cases} s_iK_{kl} & i \neq l \\ s_iK_{kl}^{-1} & i = l \end{cases}.$$

These show that SG^0 is normal in G^0 , and we may write any element of G^0 in the form $[s_1^{\epsilon_1} \dots s_n^{\epsilon_n} \phi']$, where $\epsilon_i \in \{0, 1\}$ and ϕ' is a product of extended partial conjugations and transvections. Therefore SG^0 is of index at most 2^n in G^0 . $\square$

Now let's look at the generators of $\text{Aut}(A_\Gamma)$ (respectively $\text{Out}(A_\Gamma)$) under the map $\Phi : \text{Aut}(A_\Gamma) \rightarrow \text{GL}_n(\mathbb{Z})$ (respectively $\bar{\Phi} : \text{Out}(A_\Gamma) \rightarrow \text{GL}_n(\mathbb{Z})$). If α is a graph symmetry, then $\Phi(\alpha)$ is the appropriate permutation matrix corresponding to the permutation α induces on the vertices. For a partial conjugation K_{ij} we see that $\Phi(K_{ij})$ is the identity matrix I ; Φ sends the inversion s_i to the matrix S_i which has 1 everywhere on the diagonal except for -1 at the (i, i) th position, and zeroes everywhere else, and Φ sends the transvection ρ_{ij} to the matrix $T_{ji} = I + E_{ji}$, where E_{ji} is the elementary matrix with 1 in the (i, j) th position, and zeroes everywhere else. The swapping between ρ_{ij} and T_{ji} may seem a little unnatural, but occurs as a choice of having $\text{Aut}(A_\Gamma)$ act on the left. It follows that the image of $\text{SAut}^0(A_\Gamma)$ under Φ is the subgroup of $\text{GL}_n(\mathbb{Z})$ generated by matrices of the form T_{ij} , where $v_j \leq v_i$. After restricting our attention to G^0 and SG^0 our main tool for studying the image of $\text{Out}(A_\Gamma)$ and its subgroups under $\bar{\Phi}$ will be by ordering the vertices of Γ in the manner described in Sections 2.2 and 2.3. If we order the vertices as in Section 2.2, then $[\rho_{ij}] \in \text{Out}(A_\Gamma)$ only if either v_i and v_j are in the same equivalence class of vertices, or $i \leq j$. It follows that a matrix in the image of $\bar{\Phi}|_{\text{Out}^0(A_\Gamma)}$ has a block decomposition of the form:

$$M = \begin{pmatrix} M_1 & 0 & \dots & 0 \\ * & M_2 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ * & * & \dots & M_r \end{pmatrix},$$

where the $*$ in the (i, j) th entry in the block decomposition may be nonzero if $[v_{m_j}] \leq [v_{m_i}]$, but zero otherwise.

Similarly, given a subgroup $G \leq \text{Out}(A_\Gamma)$ generated by a subset of $\mathcal{A}_\Gamma$, if we order the vertices by the method given in Section 2.3, a matrix in the image of $\bar{\Phi}|_{G^0}$ has a block decomposition of the form:

$$M = \begin{pmatrix} N_1 & 0 & \dots & 0 \\ * & N_2 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ * & * & \dots & N_{r'} \end{pmatrix},$$

where the $*$ in the (i, j) th entry in the block decomposition may be nonzero if $[v_{l_j}]_G \leq [v_{l_i}]_G$, but zero otherwise. If $[v]_G$ is an abelian equivalence class of vertices, then G contains a copy $\text{SL}(A_{[v]_G})$ generated by the $[\rho_{ij}]$ with $v_i, v_j \in [v]$. This fact lends itself to the following definition:

Definition 5.2. For any subgroup $G \leq \text{Out}(A_\Gamma)$, the SL-dimension of G , $d_{SL}(G)$, is defined to be the size of the largest abelian equivalence class under $\sim_G$.

Roughly speaking, $d_{SL}(G)$ is the largest integer such that G contains an obvious copy of $\text{SL}_{d_{SL}(G)}(\mathbb{Z})$. Note that $d_{SL}(\text{Out}(A_\Gamma))$ is simply the size of the largest abelian equivalence class under the relation $\leq$ defined on the vertices. As each

abelian equivalence class of vertices is a clique in Γ , the SL-dimension of $\text{Out}(A_\Gamma)$ is less than or equal to the size of a maximal clique in Γ (this is known as the *dimension of A_Γ*). We can now look at how G and its SL-dimension behave under restriction, exclusion, and projection maps.

Lemma 5.3. *Suppose that Γ' is a full subgraph of Γ and the conjugacy class of $A_{\Gamma'}$ in A_Γ is preserved by G . Then under the restriction map $R_{\Gamma'}$, the group $R_{\Gamma'}(G)$ is generated by a subset of $\mathcal{A}_{\Gamma'}$, and $d_{SL}(R_{\Gamma'}(G)) \leq d_{SL}(G)$.*

Proof. One first checks that for an element $[\phi] \in T$, either $R_{\Gamma'}([\phi])$ is trivial or $R_{\Gamma'}([\phi]) \in \mathcal{A}_{\Gamma'}$. This is obvious in the case of graph symmetries, inversions, and transvections. In the case of partial conjugations if v_j is not in Γ' , or if $\Gamma_{ij} \cap \Gamma' = \emptyset$, then $R_{\Gamma'}([K_{ij}])$ is trivial. Otherwise, $\Gamma_{ij} \cap \Gamma'$ is a union of connected components of $\Gamma' - st(v_j)$, so that $R_{\Gamma'}([K_{ij}])$ is an extended partial conjugation of $A_{\Gamma'}$. This proves the first part of the lemma. To prove the second part of the lemma, we first give an alternate definition of $d_{SL}(G)$. Elements in the image of G^0 under $\overline{\Phi}$ are of the form:

$$(2) \quad M = \begin{pmatrix} N_1 & 0 & \dots & 0 \\ * & N_2 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ * & * & \dots & N_{r'} \end{pmatrix},$$

where each N_i is an invertible matrix of size $l_{i+1} - l_i$. Each of the blocks is associated to either an abelian or non-abelian equivalence class in $\sim_G$, so $d_{SL}(G)$ is the size of the largest diagonal block in this decomposition associated to an abelian equivalence class. Abelian equivalence classes of $\sim_G$ are mapped to abelian equivalence classes of $\sim_{R_{\Gamma'}(G)}$, and the action of $R_{\Gamma'}(G)^0$ on $A_{\Gamma'}^{ab}$ is obtained by removing rows and columns from the decomposition given in Equation (2). Therefore, the largest diagonal block in the action of $R_{\Gamma'}(G)^0$ on $A_{\Gamma'}^{ab}$ associated to an abelian equivalence class will be of size less than or equal to $d_{SL}(G)$. Therefore $d_{SL}(R_{\Gamma'}(G)) \leq d_{SL}(G)$. $\square$

The following lemma is shown in the same way:

Lemma 5.4. *Let Γ' be a full subgraph of Γ . Suppose that the normal subgroup generated by $A_{\Gamma'}$ in A_Γ is preserved by G . Then under the exclusion map $E_{\Gamma'}$, the group $E_{\Gamma'}(G)$ is generated by a subset of $\mathcal{A}_{\Gamma'}$, and $d_{SL}(E_{\Gamma'}(G)) \leq d_{SL}(G)$.*

As projection maps are obtained by the concatenation of a restriction and an exclusion map, combining the previous two lemmas gives the following:

Proposition 5.5. *Suppose that Γ is connected and v is a maximal vertex of Γ . Under the projection homomorphism P_v of Example 2.2, the group $P_v(G^0)$ is generated by a subset of $\mathcal{A}_{lk[v]}$ and $d_{SL}(P_v(G^0)) \leq d_{SL}(G) = d_{SL}(G^0)$.*

6. PROOF OF THEOREM 6.3

Suppose that a group Λ admits no surjective homomorphisms to $\mathbb{Z}$, so that $\text{Hom}(\Lambda, \mathbb{Z}) = 0$. If a group H satisfies the property:

For every subgroup $H' \leq H$ there exists a surjective homomorphism $H' \rightarrow \mathbb{Z}$.

Then $\text{Hom}(\Lambda, H) = 0$, also. For instance, a simple induction argument on nilpotency class gives the following:

Proposition 6.1. *Suppose that $\text{Hom}(\Lambda, \mathbb{Z}) = 0$ and G is a finitely generated torsion-free nilpotent group. Then every homomorphism $f : \Lambda \rightarrow G$ is trivial.*

By Theorem 4.8 we know that $\overline{\mathcal{T}}(A_\Gamma)$ is residually torsion-free nilpotent. Combining this with Proposition 6.1 gives:

Proposition 6.2. *Suppose that $\text{Hom}(\Lambda, \mathbb{Z}) = 0$ and $f : \Lambda \rightarrow \overline{\mathcal{T}}(A_\Gamma)$ is a homomorphism. Then f is trivial.*

The overriding theme here is that we may build homomorphism rigidity results from weaker criteria by carefully studying a group's subgroups and quotients. This is also the flavour of our main theorem:

Theorem 6.3. *Suppose that G is a subgroup of $\text{Out}(A_\Gamma)$ generated by a subset $T \subset A_\Gamma$ and $d_{SL}(G) \leq m$. Let $F(\Gamma)$ be the size of a maximal, discrete, full subgraph of A_Γ . Let Λ be a group. Suppose that for each finite index subgroup $\Lambda' \leq \Lambda$, we have:*

- *Every homomorphism $\Lambda' \rightarrow \text{SL}_m(\mathbb{Z})$ has finite image,*
- *For all $N \leq F(\Gamma)$, every homomorphism $\Lambda' \rightarrow \text{Out}(F_N)$ has finite image.*
- *$\text{Hom}(\Lambda', \mathbb{Z}) = 0$*

Then every homomorphism $f : \Lambda \rightarrow G$ has finite image.

The remainder of this section will be dedicated to a proof of Theorem 6.3. We proceed by induction on the number of vertices in Γ . If Γ contains only one vertex, then $\text{Out}(A_\Gamma) \cong \mathbb{Z}/2\mathbb{Z}$, so there is no work to do. As the conditions on Λ are also satisfied by finite index subgroups, we shall allow ourselves to pass to such subgroups without further comment.

Remark 6.4. If either $m \geq 2$ or $F(\Gamma) \geq 2$, then as there exist no homomorphisms from Λ' to $\text{SL}_m(\mathbb{Z})$ or $\text{Out}(F_{F(\Gamma)})$ with infinite image, it follows that $\text{Hom}(\Lambda', \mathbb{Z}) = 0$ also. This is always the case when $G = \text{Out}(A_\Gamma)$ and $|V(\Gamma)| \geq 2$. Hence the above statement of Theorem 6.3 is a strengthening of the version given in the introduction.

Let $f : \Lambda \rightarrow G$ be such a homomorphism. There are three cases to consider:

6.1. Γ is disconnected. In this case $A_\Gamma \cong F_N *_{i=1}^k A_{\Gamma_i}$, where each Γ_i is a connected graph containing at least two vertices. Let $\Lambda' = f^{-1}(\text{Out}^0(A_\Gamma))$. As $\text{Out}^0(A_\Gamma)$ is finite index in $\text{Out}(A_\Gamma)$, this means Λ' is finite index in Λ . As in Example 2.3, for each Γ_i there is a restriction homomorphism:

$$R_i : \text{Out}^0(A_\Gamma) \rightarrow \text{Out}^0(A_{\Gamma_i}).$$

By Lemma 5.3, $R_i(G)$ is generated by a subset $T_i \subset A_{\Gamma_i}$, and $d_{SL}(R_i(G)) \leq d_{SL}(G)$. As Γ_i is a proper subgraph of Γ , we have $F(\Gamma_i) \leq F(\Gamma)$ and $|V(\Gamma_i)| < |V(\Gamma)|$. Hence, by induction $R_i f(\Lambda')$ is finite for each i , and there exists a finite index subgroup Λ_i of Λ' such that $R_i f(\Lambda_i)$ is trivial. We may also consider the exclusion homomorphism:

$$E : \text{Out}(A_\Gamma) \rightarrow \text{Out}(F_N).$$

As $N \leq F(\Gamma)$, the group $\ker(Ef)$ is a finite index subgroup of Λ . Let

$$\Lambda'' = \cap_{i=1}^k \Lambda_i \cap \ker(Ef).$$

As Λ'' is the intersection of a finite number of finite index subgroups, it is also finite index in Λ . We now study the action of Λ'' on $H_1(A_\Gamma)$. The transvection $[\rho_{ij}]$ belongs to $\text{Out}^0(A_\Gamma)$ only if either v_i and v_j belong to the same connected component of Γ , or if v_i is an isolated vertex of Γ . Therefore the action of $\text{Out}^0(A_\Gamma)$ on $H_1(A_\Gamma)$ has a block decomposition of the following form:

$$\begin{pmatrix} M_1 & 0 & \dots & 0 & * \\ 0 & M_2 & \dots & 0 & * \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & M_k & * \\ 0 & 0 & \dots & 0 & M_{k+1} \end{pmatrix},$$

where M_i corresponds to the action on A_{Γ_i} , and M_{k+1} corresponds to the action on F_N . However, as $R_i f(\Lambda'')$ is trivial for each i , and $E f(\Lambda'')$ is trivial, the action of Λ'' on $H_1(A_\Gamma)$ is of the form:

$$\begin{pmatrix} I & 0 & \dots & 0 & * \\ 0 & I & \dots & 0 & * \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & I & * \\ 0 & 0 & \dots & 0 & I \end{pmatrix}.$$

This means there is a homomorphism from Λ'' to an abelian subgroup of $\mathrm{GL}_n(\mathbb{Z})$. As $\mathrm{Hom}(\Lambda'', \mathbb{Z}) = 0$, this homomorphism must be trivial. Hence $f(\Lambda'') \subset \overline{\mathcal{T}}(A_\Gamma)$. By Proposition 6.2, this shows that $f(\Lambda'')$ is trivial. Hence $f(\Lambda)$ is finite.

6.2. Γ is connected and $Z(A_\Gamma)$ is trivial. Let $\Lambda' = f^{-1}(\mathrm{Out}^0(A_\Gamma)) = f^{-1}(G^0)$. For each maximal vertex v of Γ we have a projection homomorphism:

$$P_v : \mathrm{Out}^0(A_\Gamma) \rightarrow \mathrm{Out}^0(A_{lk[v]}).$$

By Proposition 5.5, $P_v(G^0)$ is generated by a subset $T_v \subset \mathcal{A}_{lk[v]}$ and $d_{SL}(P_v(G^0)) \leq d_{SL}(G^0) = d_{SL}(G)$. As $lk[v]$ is a proper subgraph of Γ , we have $F(lk[v]) \leq F(\Gamma)$ and $|V(lk[v])| < |V(\Gamma)|$. Therefore by induction $P_v f(\Lambda')$ is finite. Let

$$\Lambda'' = \bigcap_{[v] \text{ max.}} \ker(P_v f).$$

Then Λ'' is a finite index subgroup of Λ and lies in the kernel of the amalgamated projection homomorphism:

$$P : \mathrm{Out}^0(A_\Gamma) \rightarrow \bigoplus_{[v] \text{ max.}} \mathrm{Out}^0(A_{lk[v]}).$$

By Theorem 2.4, $\ker P$ is a finitely generated free-abelian group. As $\mathrm{Hom}(\Lambda'', \mathbb{Z}) = 0$, a homomorphism from Λ'' to $\ker P$ must be trivial. Therefore $f(\Lambda'')$ is trivial and $f(\Lambda)$ is finite.

6.3. Γ is connected and $Z(A_\Gamma)$ is nontrivial. Suppose that $Z(A_\Gamma)$ is nontrivial. Let $[v]$ be the unique maximal equivalence class in Γ . Let P_v and R_v be the restriction and projection maps given in Theorem 2.5. If $[v]$ is not equal to the whole of Γ then by induction $P_v f(\Lambda)$ and $R_v f(\Lambda)$ are both finite, therefore there exists a finite index subgroup Λ' of Λ such that $f(\Lambda')$ is contained in the kernel of $P_v \times R_v$. This is the free abelian group Tr , so the image of Λ' in Tr is trivial, and $f(\Lambda)$ is finite.

Therefore we may assume that $\Gamma = [v]$. We now look at the $\sim_G$ equivalence classes in Γ . As A_Γ is free abelian, each $[v_i]_G \subset [v]$ is abelian, and as $d_{SL}(G) \leq m$, every such $[v_i]_G$ contains at most m vertices. Therefore matrices in (the image of) G^0 (under Φ) are of the form:

$$M = \begin{pmatrix} N_1 & 0 & \dots & 0 \\ * & N_2 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ * & * & \dots & N_{r'} \end{pmatrix},$$

where the $*$ in the (i, j) th block is possibly nonzero if $[v_i] \leq [v_j]$. For each i , we can look at the projection $M \mapsto N_i$ to obtain a homomorphism $g_i : SG^0 \rightarrow \mathrm{SL}_{l_{i+1}-l_i}(\mathbb{Z})$. As $l_{i+1} - l_i \leq m$, our hypothesis on Λ implies that $g_i f(f^{-1}(SG^0))$ is finite for all i . Let Λ_i be the kernel of each map $g_i f$ restricted to $f^{-1}(SG^0)$. Each

Λ_i is finite index in Λ . Let $\Lambda' = \cap_{i=1}^k \Lambda_i$. Then matrices in the image of Λ' under f are of the form:

$$M = \begin{pmatrix} I & 0 & \dots & 0 \\ * & I & \dots & 0 \\ \dots & \dots & \dots & \dots \\ * & * & \dots & I \end{pmatrix},$$

therefore $f(\Lambda')$ is a torsion-free nilpotent group. By Proposition 6.1, $f(\Lambda')$ is trivial. Hence $f(\Lambda)$ is finite, and this finishes the final case of the theorem.

7. CONSEQUENCES OF THEOREM 6.3

We say that a group Λ is $\mathbb{Z}$ -averse if no finite index subgroup of Λ contains a normal subgroup that maps surjectively to $\mathbb{Z}$. This restriction gives a large class of groups to which Λ cannot map. For instance, in [6] Bridson and the author prove the following theorem:

Theorem 7.1. *Suppose that Λ is $\mathbb{Z}$ -averse and $f : \Lambda \rightarrow \text{Out}(F_n)$ is a homomorphism. Then $f(\Lambda)$ is finite.*

Note that if Λ is $\mathbb{Z}$ -averse, then every finite index subgroup of Λ is also $\mathbb{Z}$ -averse. As there are no homomorphisms from a $\mathbb{Z}$ -averse group to $\text{SL}_2(\mathbb{Z})$ with infinite image (as $\text{SL}_2(\mathbb{Z})$ is virtually free), combining the above with Theorem 6.3 we obtain:

Corollary 7.2. *If Λ is a $\mathbb{Z}$ -averse group, and Γ is a finite graph that satisfies $d_{SL}(\text{Out}(A_\Gamma)) \leq 2$, then every homomorphism $f : \Lambda \rightarrow \text{Out}(A_\Gamma)$ has finite image.*

We would like to apply Theorem 6.3 to higher-rank lattices in Lie groups. For the remainder of this section Λ will be an irreducible lattice in a semisimple real Lie group G with real rank $\text{rank}_{\mathbb{R}} G \geq 2$, finite centre, and no compact factors. Such lattices are $\mathbb{Z}$ -averse by Margulis' normal subgroup theorem, which states that if Λ' is a normal subgroup of Λ then either Λ/Λ' is finite, or $\Lambda' \subset Z(G)$, so Λ' is finite. The work of Margulis also lets us restrict the linear representations of such lattices:

Proposition 7.3. *If $\text{rank}_{\mathbb{R}} G \geq k$ then every homomorphism $f : \Lambda \rightarrow \text{SL}_k(\mathbb{Z})$ has finite image.*

To prove this we appeal to Margulis superrigidity. The following two theorems follow from [19], Chapter IX, Theorems 6.15 and 6.16 and the remarks in 6.17:

Theorem 7.4. *Let H be a real algebraic group and $f : \Lambda \rightarrow H$ a homomorphism. The Zariski closure of the image of f , denoted $\overline{f(\Lambda)}$, is semisimple.*

Theorem 7.5 (Margulis' Superrigidity Theorem). *Let H be a connected, semisimple, real algebraic group and $f : \Lambda \rightarrow H$ a homomorphism. If*

- *H is adjoint (equivalently $Z(H) = 1$) and has no compact factors, and*
- *$f(\Lambda)$ is Zariski dense in H ,*

then f extends uniquely to a continuous homomorphism $\tilde{f} : G \rightarrow H$. Furthermore, if $Z(G) = 1$ and $f(\Lambda)$ is nontrivial and discrete, then $\tilde{f}$ is an isomorphism.

We may combine these to prove Proposition 7.3:

Proof of Proposition 7.3. Let $f : \Lambda \rightarrow \text{SL}_k(\mathbb{Z})$ be a homomorphism. By Theorem 7.4, the Zariski closure of the image $\overline{f(\Lambda)} \subset \text{SL}_k(\mathbb{R})$ is semisimple. Also, $\overline{f(\Lambda)}$ has finitely many connected components — let $\overline{f(\Lambda)}_0$ be the connected component containing the identity. Decompose $\overline{f(\Lambda)}_0 = H_1 \times K$, where K is a maximal compact factor. Then H_1 is a connected semisimple real algebraic group with no compact factors. We look at the finite index subgroup $\Lambda_1 = f^{-1}(H_1)$ of Λ , so that

$\overline{f(\Lambda_1)} = H_1$. As the centre of a subgroup of an algebraic group is contained in the centre of its Zariski closure, $f(Z(\Lambda_1)) \subset Z(H_1)$. This allows us to factor out centres in the groups involved. Let $G_2 = G/Z(G)$, $\Lambda_2 = \Lambda_1/Z(\Lambda_1) = \Lambda_1/(\Lambda_1 \cap Z(G))$ and $H_2 = H_1/Z(H_1)$. Then there is an induced map $f_2 : \Lambda_2 \rightarrow H_2$ satisfying the conditions of Theorem 7.5. Therefore if $f_2(\Lambda_2) \neq 1$ there is an isomorphism $\tilde{f}_2 : G_2 \rightarrow H_2$. However

$$\begin{aligned} \text{rank}_{\mathbb{R}} G_2 &= \text{rank}_{\mathbb{R}} G \geq k \\ \text{rank}_{\mathbb{R}} H_2 &= \text{rank}_{\mathbb{R}} H_1 \leq \text{rank}_{\mathbb{R}} \text{SL}_k(\mathbb{R}) = k - 1. \end{aligned}$$

This contradicts the isomorphism between H_2 and G_2 . Therefore $f_2(\Lambda_2) = 1$. As $Z(\Lambda_1)$ is finite, and Λ_1 is finite index in Λ , this shows that the image of Λ under f is finite. $\square$

Combining Proposition 7.3 with Theorems 7.1 and 6.3, this gives:

Theorem 7.6. *Let G be a real semisimple Lie group with finite centre, no compact factors, and $\text{rank}_{\mathbb{R}} G \geq 2$. Let Λ be an irreducible lattice in G . If $\text{rank}_{\mathbb{R}} G \geq d_{\text{SL}}(\text{Out}(A_{\Gamma}))$, then every homomorphism $f : \Lambda \rightarrow \text{Out}(A_{\Gamma})$ has finite image.*

The following corollary justifies our definition of SL-dimension, and shows that you can't hide any larger copies of SL_n inside $\text{Out}(A_{\Gamma})$:

Corollary 7.7. *$\text{Out}(A_{\Gamma})$ contains a subgroup isomorphic to $\text{SL}_k(\mathbb{Z})$ if and only if $k \leq d_{\text{SL}}(\text{Out}(A_{\Gamma}))$.*

REFERENCES

- [1] Hyman Bass and Alexander Lubotzky. Linear-central filtrations on groups. In *The mathematical legacy of Wilhelm Magnus: groups, geometry and special functions (Brooklyn, NY, 1992)*, volume 169 of *Contemp. Math.*, pages 45–98. Amer. Math. Soc., Providence, RI, 1994.
- [2] M. Bestvina and N. Brady. Morse theory and finiteness properties for groups, *Invent. Math.* 129 (1997), 445–470.
- [3] M. Bestvina and M. Feighn. A hyperbolic $\text{Out}(F_n)$ -complex, *arXiv:0808.3730*, 2009.
- [4] Nicolas Bourbaki. *Lie groups and Lie algebras. Chapters 1–3*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 1989.
- [5] Martin R. Bridson and Benson Farb. A remark about actions of lattices on free groups. *Topology Appl.*, 110(1):21–24, 2001.
- [6] Martin R. Bridson and Richard D. Wade. Actions of higher-rank lattices on free groups. *arXiv:1004.3222*, 2010.
- [7] Ruth Charney. An introduction to right-angled Artin groups. *Geom. Dedicata*, 125:141–158, 2007.
- [8] Ruth Charney and Karen Vogtmann. Finiteness properties of automorphism groups of right-angled Artin groups. *Bull. Lond. Math. Soc.*, 41(1):94–102, 2009.
- [9] Ruth Charney and Karen Vogtmann. Subgroups and quotients of automorphism groups of RAAGs. *arXiv:0909.2444*, 2010.
- [10] F. Dahmani, V. Guirardel and D. Osin. Hyperbolic embeddings and rotating families in groups acting on hyperbolic spaces, in preparation: preliminary version www.math.univ-toulouse.fr/~dahmani/Files/VRF.pdf.
- [11] Matthew B. Day. Symplectic structures on right-angled Artin groups: between the mapping class group and the symplectic group. *Geom. Topol.*, 13(2):857–899, 2009.
- [12] G. Duchamp and D. Krob. The free partially commutative Lie algebra: bases and ranks. *Adv. Math.*, 95(1):92–126, 1992.
- [13] G. Duchamp and D. Krob. The lower central series of the free partially commutative group. *Semigroup Forum*, 45(3):385–394, 1992.
- [14] Benson Farb and Howard Masur. Superrigidity and mapping class groups. *Topology*, 37(6):1169–1176, 1998.
- [15] Frédéric Haglund and Daniel T. Wise. Special cube complexes. *Geom. Funct. Anal.*, 17(5):1551–1620, 2008.
- [16] M. Handel and L. Mosher. Subgroup classification in $\text{Out}(F_n)$. *arXiv:0908.1255*, 2009.
- [17] Michael R. Laurence. A generating set for the automorphism group of a graph group. *J. London Math. Soc. (2)*, 52(2):318–334, 1995.

- [18] Roger C. Lyndon and Paul E. Schupp. *Combinatorial group theory*. Springer-Verlag, Berlin, 1977. *Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 89*.
- [19] Gregori Margulis *Discrete subgroups of semisimple Lie groups* Springer-Verlag, Berlin, 1991.
- [20] Wilhelm Magnus, Abraham Karrass, and Donald Solitar. *Combinatorial group theory: Presentations of groups in terms of generators and relations*. Interscience Publishers [John Wiley & Sons, Inc.], New York-London-Sydney, 1966.
- [21] Ashot Minasyan. Hereditary conjugacy separability of right angled Artin groups and its applications. *arXiv:0905.1282*, 2009.
- [22] Alexandra Pettet. The Johnson homomorphism and the second cohomology of IA_n . *Algebr. Geom. Topol.*, 5:725–740, 2005.
- [23] Herman Servatius. Automorphisms of graph groups. *J. Algebra*, 126(1):34–60, 1989.
- [24] Takao Satoh. New obstructions for the surjectivity of the Johnson homomorphism of the automorphism group of a free group. *J. London Math. Soc. (2)*, 74(2):341–360, 2006.
- [25] Emmanuel Toinet. Conjugacy p -separability of right-angled Artin groups and applications. *arXiv:1009.3859*, 2010.

MATHEMATICAL INSTITUTE, 24-29 ST GILES', OXFORD OX1 3LB, UK
E-mail address: `wade@maths.ox.ac.uk`